

# Gamma and X-Ray Standards




A wide range of gamma and x-ray standards for research and educational use are available in the energy range of 5.9 to 2614 keV. Many nuclides are available up to 100  $\mu$ Ci (3.7 MBq) to allow for the lower efficiencies of detection inherent in many instruments. Available activity ranges are listed for each nuclide in the tables on pages 46 to 53. Please contact customer service for other activity requirements.

## Gamma Standards—Type C

The type C configuration can be used to check the performance of G.M. and NaI (TI) detectors. The maximum activity of this source type is 10  $\mu$ Ci (370 kBq). Please call customer service for a quotation.

**Figure 45-A: Type C**

**A1002**



### Overall Dimensions

| Height            | Width          | Length            | Active Diameter |
|-------------------|----------------|-------------------|-----------------|
| 0.075"<br>1.91 mm | 0.43"<br>11 mm | 0.925"<br>23.5 mm | 0.118"<br>3 mm  |

### Window & Active Deposit

| Window  | Nature of Active Deposit  |
|---------|---------------------------|
| Plastic | Evaporated Metallic Salts |

### Simulated I-125 Sources

I-129 sources are frequently referred to as "simulated I-125 sources." I-129 does not in fact simulate I-125 well enough to use it to calibrate an I-125 counter. The equivalence of I-129 to I-125 will vary from one counter to another due to the differing photon energies and counting geometries. EZIP recommends the purchase of an I-125 standard for the initial instrument calibration. An I-129 source may then be used for a daily check of instrument response.

The energies and abundances of I-125 and I-129 x-rays and photons are:

|       | K x-rays (keV)     | Gammas (keV) |
|-------|--------------------|--------------|
| I-125 | (Te) 27-32 (141%)  | 35.5 (6.66%) |
| I-129 | (Xe) 29-35 (70.8%) | 39.6 (7.5%)  |


# Gamma and X-Ray Standards

## Gamma Standards—Type D

The type D configuration is mainly used for checking the performance of G.M. and NaI(Tl) detectors. The type D disk is a 1" (25.4 mm) diameter by 0.250" (6.35 mm) thick disk constructed of high strength plastic. The active diameter is 0.197" (5 mm).



**Figure 46-A: Type D Disk**



| Overall Dimensions |                 |         |
|--------------------|-----------------|---------|
| Overall Diameter   | Active Diameter | Height  |
| 1"                 | 0.197"          | 0.25"   |
| 25.4 mm            | 5 mm            | 6.35 mm |

| Window & Active Deposit |                           |
|-------------------------|---------------------------|
| Window                  | Nature of Active Deposit  |
| Plastic                 | Evaporated Metallic Salts |

**Gamma Standards—Type D**

| Catalog Number | Nuclide       | Half-Life                | Major Photon Emissions (keV)  | Available Activities |                 |
|----------------|---------------|--------------------------|---|----------------------|-----------------|
| GF-241-D       | Americium-241 | 432.17 y                 | 59.5 (36%), 11-20 (39.5%) Np L x-rays   | 5 nCi-100 µCi        | 185 Bq-3.7 MBq  |
| GF-124-D       | Antimony-124  | 60.20 d                  | 602.7 (97.9%), 722.8 (10.9%), 1690.9 (47.6%)  | 5 nCi-100 µCi        | 185 Bq-3.7 MBq  |
| GF-125A-D      | Antimony-125  | 1007.7 d                 | 428 (29.7%), 464 (10.5%), 601 (17.7%), 607 (5%), 636 (11.2%), 27-32 (75.1%) Te K x-rays | 5 nCi-100 µCi        | 185 Bq-3.7 MBq  |
| GF-133-D       | Barium-133    | 3862 d                   | 80 (34.1%), 303 (18.3%), 356 (61.9%), 32-37 (53.2%) Cs K x-rays                         | 5 nCi-100 µCi        | 185 Bq-3.7 MBq  |
| GF-007-D       | Beryllium-7   | 53.284 d                 | 478 (10.3%)   | 25 nCi-500 µCi       | 925 Bq-18.5 MBq |
| GF-207-D       | Bismuth-207   | 1.16 x 10 <sup>4</sup> d | 570 (97.7%), 1064 (74.5%), 9-15 (32.5%), Pb L x-rays, 72-88 (77.7%) Pb K x-rays         | 5 nCi-100 µCi        | 185 Bq-3.7 MBq  |
| GF-109-D       | Cadmium-109   | 462.6 d                  | 88 (3.6% from Ag-109m), 22-26 (99.4%) Ag K x-rays                                       | 50 nCi-1 mCi         | 1.85 kBq-37 MBq |
| GF-139-D       | Cerium-139    | 137.640 d                | 33.03 (22.8%), 33.4 (41.9%), 165.9 (79.9%), 33-39 (80%) La x-rays                       | 5 nCi-10 µCi         | 185 Bq-370 kBq  |
| GF-141-D       | Cerium-141    | On Request               | —   | —                    | —               |
| GF-134-D       | Cesium-134    | 754.28 d                 | 563 (8.4%), 569 (15.4%), 605 (97.6%), 796 (85.4%)                                       | 5 nCi-100 µCi        | 185 Bq-3.7 MBq  |
| GF-137-D       | Cesium-137    | 30.17 y                  | 662 (85.1% from Ba-137), 32-37 (7.2%) Ba K x-rays                                       | 5 nCi-1 mCi          | 185 Bq-3.7 MBq  |
| GF-051-D       | Chromium-51   | 27.706 d                 | 320 (9.86%), 4.9-5.4 (22.8%) V K x-rays   | 25 nCi-500 µCi       | 185 Bq-37 MBq   |

| Gamma Standards—Type D |                                  |                           |  |                      |                  |  |
|------------------------|----------------------------------|---------------------------|--|----------------------|------------------|--|
| Catalog Number         | Nuclide                          | Half-Life                 | Major Photon Emissions (keV)   | Available Activities |                  |  |
| GF-056-D               | Cobalt-56                        | 77.31 d                   | 846.8 (99.9%), 1238 (66.1%), 1771 (15.5%), 2035 (7.8%), 2598 (17%), 3253 (7.6%), others up to 3452 | 10 nCi-100 µCi       | 370 Bq-3.7 MBq   |  |
| GF-057-D               | Cobalt-57                        | 271.79 d                  | 14 (9.2%), 122 (85.6%), 136.5 (10.7%), 6.4-7.1 (57.9%) Fe K x-rays                                 | 5 nCi-1 mCi          | 185 Bq-37 MBq    |  |
| GF-058-D               | Cobalt-58                        | 70.86 d                   | 810 (99.5%), 6.4-7.1 (26.7%) Fe K x-rays   | 5 nCi-100 µCi        | 185 Bq-3.7 MBq   |  |
| GF-060-D               | Cobalt-60                        | 5.272 y                   | 1173 (100%), 1333 (100%)   | 5 nCi-1 mCi          | 185 Bq-37 MBq    |  |
| GF-152-D               | Europium-152                     | 4933 d                    | 122-1408, 40-47 (74%) Sm + Gd x-rays   | 25 nCi-100 µCi       | 925 Bq-3.7 MBq   |  |
| GF-154-D               | Europium-154                     | 3136.8 d                  | 123-1597, 42-50 (25.6%) Gd x-rays  | 25 nCi-100 µCi       | 925 Bq-3.7 MBq   |  |
| GF-155-D               | Europium-155                     | 1770 d                    | 87 (34%), 105 (20.6%), 42-50 (24.0%), Gd K x-rays  | 10 nCi-100 µCi       | 370 Bq-3.7 MBq   |  |
| GF-153-D               | Gadolinium-153                   | 242 d                     | 97 (29.5%), 103 (21.1%), 40-49 (122%) Eu K x-rays  | 5 nCi-1 mCi          | 185 Bq-37 MBq    |  |
| GF-068-D               | Germanium-68                     | 270.8 d                   | 511 (178%), 1077 (3.2%) from Ga-68, 9.2-10.4 (44.1%) Ga K x-rays, 8.6-9.6 (4.7%) Zn K x-rays       | 100 nCi-100 µCi      | 3.7 kBq-3.7 MBq  |  |
| GF-166-D               | Holmium-166m <sup>(1)</sup>      | 1200 y                    | 81-1427, 48-58 (37.6%) Er K x-rays   | 10 nCi-10 µCi        | 370 Bq-370 kBq   |  |
| GF-125-D               | Iodine-125                       | 59.43 d                   | 35 (6.58%), 27-32 (139%) Te K x-rays   | 10 nCi-100 µCi       | 370 Bq-3.7 MBq   |  |
| GF-129-D               | Iodine-129                       | 1.57 x 10 <sup>7</sup> y  | 40 (7.5%), 29-35 (70.4%) Xe K x-rays   | 50 nCi-1 µCi         | 1.85 kBq-37 kBq  |  |
| GF-059-D               | Iron-59                          | 44.51 d                   | 1099 (56.3%), 1292 (43.7%)   | 5 nCi-100 µCi        | 185 Bq-3.7 MBq   |  |
| GF-054-D               | Manganese-54                     | 312.3 d                   | 835 (100%), 5.4-5.9 (25.6%) Cr K x-rays  | 5 nCi-100 µCi        | 185 Bq-3.7 MBq   |  |
| GF-203-D               | Mercury-203                      | 46.595 d                  | 279.2 (81.5%)  | 10 nCi-50 µCi        | 370 Bq-1.85 MBq  |  |
| GF-226-D               | Radium-226                       | 1600 y                    | 47-2448 (includes daughters)   | 50 nCi-10 µCi        | 1.85 kBq-370 kBq |  |
| GF-106-D               | Ruthenium-106                    | 1.020 y                   | 512 (20.7%), 622 (9.8%) from Rh-106  | 25 nCi-100 µCi       | 925 Bq-3.7 MBq   |  |
| GF-046-D               | Scandium-46                      | 83.79 d                   | 889 (99.9%), 1121 (99.9%)  | 5 nCi-100 µCi        | 185 Bq-3.7 MBq   |  |
| GF-075-D               | Selenium-75                      | 119.64 d                  | 121 (17.1%), 136 (58.8%), 265 (59%), 280 (25%), 10.5-12.0 (56.8%) As K x-rays                      | 5 nCi-100 µCi        | 185 Bq-3.7 MBq   |  |
| GF-110-D               | Silver-110m                      | 249.8 d                   | 657.8 (94.4%), 884.6 (72.6%)   | 5 nCi-50 µCi         | 185 Bq-1.85 MBq  |  |
| GF-131-D               | Simulated I-131                  | ~5 y                      | 356 (from Ba-133), 662 (from Cs-137/Ba-137)  | 50 nCi-100 µCi       | 1.85 kBq-3.7 MBq |  |
| GF-022-D               | Sodium-22                        | 950.8 d                   | 511 (178%), 1275 (100%)  | 5 nCi-100 µCi        | 185 Bq-3.7 MBq   |  |
| GF-085-D               | Strontium-85                     | 64.849 d                  | 514 (98.4%), 13.3-15.3 (58.7%) Rb K x-rays   | 5 nCi-100 µCi        | 185 Bq-3.7 MBq   |  |
| GF-228-D               | Thorium-228                      | 698.2 d                   | 84-2614 (includes daughters)   | 10 nCi-10 µCi        | 370 Bq-370 kBq   |  |
| GF-113-D               | Tin-113                          | 115.09 d                  | 392 (64% from In-113 m), 24-28 (96.8%) In K x-rays   | 5 nCi-100 µCi        | 185 Bq-3.7 MBq   |  |
| GF-235-D               | Uranium-235 <sup>(1)</sup>       | 7.037 x 10 <sup>8</sup> y | 143 (10.5%), 186 (53%), 90-105 (10.8%) Th K x-rays   | 10 nCi-100 nCi       | 370 Bq-3.7 kBq   |  |
| GF-238U-D              | Uranium (Natural) <sup>(1)</sup> | 4.468 x 10 <sup>9</sup> y | 26-2448 (includes daughters)   | 10 nCi-100 nCi       | 370 Bq-3.7 kBq   |  |
| GF-088-D               | Yttrium-88                       | 106.630 d                 | 898 (94%), 1836 (99.4%), 14.1-16.2 (61.6%) Sr K x-rays   | 5 nCi-100 µCi        | 185 Bq-3.7 MBq   |  |
| GF-065-D               | Zinc-65                          | 244.26 d                  | 1116 (50.6%), 8.0-8.9 (38.7%) Cu K x-rays  | 5 nCi-100 µCi        | 185 Bq-3.7 MBq   |  |
| GF-095-D               | Zirconium-95/Nb-95               | 64.02 d                   | 724 (44.1%), 757 (54.5%)   | 10 nCi-50 µCi        | 370 Bq-1.85 MBq  |  |

1) 9 mm active diameter.

| Gamma Sets—Type D |                      |                 |   |
|-------------------|----------------------|-----------------|---|
| Catalog Number    | Available Activities | Sets Consist Of |   |
| GF-290-0.1D       | 0.1 µCi              | 3.7 kBq         | Ba-133, Cd-109, Co-57, Co-60, Cs-137, Mn-54 and Na-22 |
| GF-290-1D         | 1 µCi                | 37 kBq          | Ba-133, Cd-109, Co-57, Co-60, Cs-137, Mn-54 and Na-22 |
| GF-290-10D        | 10 µCi               | 370 kBq         | Ba-133, Cd-109, Co-57, Co-60, Cs-137, Mn-54 and Na-22 |

# Gamma and X-Ray Standards


## Gamma Standards—Type M

The type M thin "scatterless" disk is used in applications involving high resolution solid state detectors. The activity is deposited on 9 mg/cm<sup>2</sup> aluminized Mylar (polyester) disk, and covered with 0.9 mg/cm<sup>2</sup> Kapton (polyimide). The source is supplied in a removable aluminum holder. In the holder the source has an overall diameter of 1" (25.4 mm) and a thickness of 0.125" (3.18 mm). Out of the holder the source is 0.937" (23.8 mm) in diameter with a thickness of approximately 0.030" (0.76 mm). The active diameter is 0.12" (3 mm).



**Figure 48-A: Type M Disk**

A1202



### Overall Dimensions

| Overall Diameter | Active Diameter | Height  |
|------------------|-----------------|---------|
| 1"               | 0.12"           | 0.125"  |
| 25.4 mm          | 3 mm            | 3.18 mm |

| Window   | Nature of Active Deposit    |
|--|-----------------------------|
| 0.010" (0.254 mm)                                      | Evaporated Aluminized Mylar |
| Metallic Salts   |                             |
| <b>Exceptions</b>                                      |                             |
| Fe-55: 0.00025" (0.0064 mm) Aluminized Mylar           |                             |
| Am-241: 0.001" (0.0254 mm) Aluminum and Platinum Foils |                             |

**Gamma Standards—Type M**

| Catalog Number | Nuclide       | Half-Life            | Major Photon Emissions (keV)  | Available Activities |                  |
|----------------|---------------|----------------------|---|----------------------|------------------|
| GF-241-M       | Americium-241 | 432.17 y             | 59.5(36%), 11-20(39.5%) Np L x-rays   | 5 nCi-10 µCi         | 185 Bq-370 kBq   |
| GF-124-M       | Antimony-124  | 60.20 d              | 602.7(97.9%), 722.8(10.9%), 1690.9(47.6%)   | 5 nCi-100 µCi        | 185 Bq-3.7 MBq   |
| GF-125A-M      | Antimony-125A | 1007.7 d             | 428(29.7%), 464(10.5%), 601(17.7%), 607(5%), 636(11.2%), 27-32(75.1%) Te K x-rays | 5 nCi-100 µCi        | 185 Bq-3.7 MBq   |
| GF-133-M       | Barium-133    | 3862 d               | 80(34.1%), 303(18.3%), 356(61.9%), 32-37(53.2%) Cs K x-rays                       | 5 nCi-100 µCi        | 185 Bq-3.7 MBq   |
| GF-007-M       | Beryllium-7   | 53.28 d              | 478(10.3%)  | 25 nCi-500 µCi       | 925 Bq-18.5 MBq  |
| GF-207-M       | Bismuth-207   | $1.16 \times 10^4$ d | 570(97.7%), 1064(74.5%), 9-15(32.5%), Pb L x-rays, 72-88(77.7%) Pb K x-rays       | 5 nCi-100 µCi        | 185 Bq-3.7 MBq   |
| GF-109-M       | Cadmium-109   | 462.6 d              | 88(3.6% from Ag-109 m), 22-26(99.4%) Ag K x-rays                                  | 50 nCi-100 µCi       | 1.85 kBq-3.7 MBq |
| GF-139-M       | Cerium-139    | 137.640 d            | 33.03(22.8%), 33.4(41.9%), 165.9(79.9%), 33-39(80%) La x-rays                     | 50nCi-1 µCi          | 1.85 kBq-37 kBq  |
| GF-141-M       | Cerium-141    | 32.5 d               | 36.0 (9.1%), 35.6 (5%), 145.4 (48.4%), 352.42 (17%), Pr x-rays                    | On Request           | —                |
| GF-134-M       | Cesium-134    | 754.28 d             | 563(8.4%), 569(15.4%), 605(97.6%), 796(85.4%)                                     | 5 nCi-100 µCi        | 185 Bq-3.7 MBq   |
| GF-137-M       | Cesium-137    | 30.17 y              | 662(85.1% from Ba-137), 32-37(7.2%) Ba K x-rays                                   | 5 nCi-100 µCi        | 185 Bq-3.7 MBq   |
| GF-051-M       | Chromium-51   | 27.706 d             | 320(9.86%), 4.9-5.4(22.8%) V K x-rays   | 25 nCi-100 µCi       | 925 Bq-3.7 MBq   |

### Gamma Standards—Type M

| Catalog Number | Nuclide            | Half-Life | Major Photon Emissions (keV)   | Available Activities |                  |
|----------------|--------------------|-----------|--|----------------------|------------------|
| GF-056-M       | Cobalt-56          | 77.31 d   | 846.8(99.9%), 1238(66.1%), 1771(15.5%), 2035(7.8%), 2598(17%), 3253(7.6%), others up to 3452 | 10 nCi-100 µCi       | 370 Bq-3.7 MBq   |
| GF-057-M       | Cobalt-57          | 271.79 d  | 14(9.2%), 122(85.6%), 136.5(10.7%), 6.4-7.1(57.9%) Fe K x-rays                               | 5 nCi-100 µCi        | 185 Bq-3.7 MBq   |
| GF-058-M       | Cobalt-58          | 70.86 d   | 810(99.5%), 6.4-7.1(26.7%) Fe K x-rays   | 5 nCi-100 µCi        | 185 Bq-3.7 MBq   |
| GF-060-M       | Cobalt-60          | 5.272 y   | 1173(100%), 1333(100%)   | 5 nCi-100 µCi        | 185 Bq-3.7 MBq   |
| GF-152-M       | Europium-152       | 4933 d    | 122-1408, 40-47(74%) Sm +Gd x-rays   | 25 nCi-100 µCi       | 925 Bq-3.7 MBq   |
| GF-154-M       | Europium-154       | 3136.8 d  | 123-1597, 42-50(25.6%) Gd x-rays   | 25 nCi-100 µCi       | 925 Bq-3.7 MBq   |
| GF-155-M       | Europium-155       | 1770 d    | 87(34%), 105(20.6%), 42-50(24.0%), Gd K x-rays   | 10 nCi-100 µCi       | 370 Bq-3.7 MBq   |
| GF-153-M       | Gadolinium-153     | 242 d     | 97(29.5%), 103(21.1%), 40-49(122%) Eu K x-rays   | 10 nCi-100 µCi       | 370 Bq-3.7 MBq   |
| GF-068-M       | Germanium-68       | 270.8 d   | 511(178%), 1077(3.2%) from Ga-68, 9.2-10.4(44.1%) Ga K x-rays, 8.6-9.6(4.7%) Zn K x-rays     | 100 nCi-100 µCi      | 3.7 kBq-3.7 MBq  |
| GF-166-M       | Holmium-166m       | 1200 y    | 81-1427, 48-58(37.6%) Er K x-rays  | 10 nCi-1 µCi         | 370 Bq-37 kBq    |
| GF-055-M       | Iron-55            | 999 d     | 5.8-6.5(27.3%) Mn K x-rays   | 1 µCi-100 µCi        | 37 kBq-3.7 MBq   |
| GF-059-M       | Iron-59            | 44.51 d   | 1099(56.3%), 1292(43.7%)   | 5 nCi-100 µCi        | 185 Bq-3.7 MBq   |
| GF-054-M       | Manganese-54       | 312.3 d   | 835(100%), 5.4-5.9(25.6%) Cr K x-rays  | 5 nCi-100 µCi        | 185 Bq-3.7 MBq   |
| GF-203-M       | Mercury-203        | 46.595 d  | 279.2(81.5%)   | 10 nCi-10 µCi        | 370 Bq-370 kBq   |
| GF-046-M       | Scandium-46        | 83.79 d   | 889(99.9%), 1121(99.9%)  | 5 nCi-100 µCi        | 185 Bq-3.7 MBq   |
| GF-075-M       | Selenium-75        | 119.64 d  | 121(17.1%), 136(58.8%), 265(59%), 280(25%), 10.5-12.0(56.8%) As K x-rays                     | 5 nCi-100 µCi        | 185 Bq-3.7 MBq   |
| GF-110-M       | Silver-110m        | 249.8 d   | 657.8(94.4%), 884.6(72.6%)   | 5 nCi-50 µCi         | 185 Bq-1.85 MBq  |
| GF-131-M       | Simulated I-131    | ~5 y      | 356(from Ba-133), 662(from Cs-137/Ba-137)  | 50 nCi-100 µCi       | 1.85 kBq-3.7 MBq |
| GF-022-M       | Sodium-22          | 950.8 d   | 511(178%), 1275(100%)  | 5 nCi-100 µCi        | 185 Bq-3.7 MBq   |
| GF-085-M       | Strontium-85       | 64.849 d  | 514(98.4%), 13.3-15.3(58.7%) Rb K x-rays   | 5 nCi-100 µCi        | 185 Bq-3.7 MBq   |
| GF-113-M       | Tin-113            | 115.09 d  | 392(64% from In-113 m), 24-28(96.8%) In K x-rays   | 5 nCi-10 µCi         | 185 Bq-370 kBq   |
| GF-088-M       | Yttrium-88         | 106.630 d | 898(94%), 1836(99.4%), 14.1-16.2(61.6%) Sr K x-rays  | 5 nCi-100 µCi        | 185 Bq-3.7 MBq   |
| GF-065-M       | Zinc-65            | 244.26 d  | 1116(50.6%), 8.0-8.9(38.7%) Cu K x-rays  | 5 nCi-100 µCi        | 185 Bq-3.7 MBq   |
| GF-095-M       | Zirconium-95/Nb-95 | 64.02 d   | 724(44.1%), 757(54.5%)   | 10 nCi-50 µCi        | 370 Bq-1.85 MBq  |

### Gamma Sets—Type M

| Catalog Number | Available Activities | Sets Consist Of  |
|----------------|----------------------|--|
| GF-290-0.1M    | 0.1 µCi              | 3.7 kBq<br>Ba-133, Cd-109, Co-57, Co-60, Cs-137, Mn-54 and Na-22 |
| GF-290-1M      | 1 µCi                | 37 kBq<br>Ba-133, Cd-109, Co-57, Co-60, Cs-137, Mn-54 and Na-22  |
| GF-290-10M     | 10 µCi               | 370 kBq<br>Ba-133, Cd-109, Co-57, Co-60, Cs-137, Mn-54 and Na-22 |

# Gamma and X-Ray Standards

## Gamma Standards—Type R

The type R rod is used for calibrating well type NaI(Tl) detectors. It is constructed of high strength plastic and is offered in three sizes: 5" high x 0.625" diameter (127 mm x 15.9 mm), 5" high x 0.5" diameter (127 mm x 12.7 mm) and 2.95" x 0.5" diameter (74.9 mm x 12.7 mm). The active diameter of the rod standard is 0.187" (4.75 mm).




| Window & Active Materials |                            |
|---------------------------|----------------------------|
| Window                    | Nature of Active Materials |
| Plastic                   | Evaporated Metallic Salts  |

### Overall Dimensions

| Assembly | X  | Height          | Diameter         | Active Diameter  |
|----------|----|-----------------|------------------|------------------|
| A1100    | R1 | 5" (127 mm)     | 0.625" (15.9 mm) | 0.187" (4.75 mm) |
| A1102    | R2 | 5" (127 mm)     | 0.5" (12.7 mm)   | 0.187" (4.75 mm) |
| A1103    | R3 | 2.95" (74.9 mm) | 0.5" (12.7 mm)   | 0.187" (4.75 mm) |

Figure 50-A: Type R Rod



## Gamma Standards—Type R

| Catalog Number | Nuclide       | Half-Life            | Major Photon Emissions (keV)  | Available Activities |                 |
|----------------|---------------|----------------------|---|----------------------|-----------------|
| GF-241-x       | Americium-241 | 432.17 y             | 59.5 (36%), 11-20 (39.5%) Np L x-rays   | 5 nCi-100 µCi        | 185 Bq-3.7 MBq  |
| GF-124-x       | Antimony-124  | 60.20 d              | 602.7(97.9%), 722.8(10.9%), 1690.9(47.6%)   | 5 nCi-100 µCi        | 185 Bq-3.7 MBq  |
| GF-125A-x      | Antimony-125A | 1007.7 d             | 428(29.7%), 464(10.5%), 601(17.7%), 607(5%), 636(11.2%), 27-32(75.1%) Te K x-rays | 5 nCi-100 µCi        | 185 Bq-3.7 MBq  |
| GF-133-x       | Barium-133    | 3862 d               | 80(34.1%), 303(18.3%), 356(61.9%), 32-37(53.2%) Cs K x-rays                       | 5 nCi-100 µCi        | 185 Bq-3.7 MBq  |
| GF-007-x       | Beryllium-7   | On Request           |   |                      |                 |
| GF-207-x       | Bismuth-207   | $1.16 \times 10^4$ d | 570(97.7%), 1064(74.5%), 9-15(32.5%), Pb L x-rays, 72-88(77.7%) Pb K x-rays       | 5 nCi-100 µCi        | 185 Bq-3.7 MBq  |
| GF-109-x       | Cadmium-109   | 462.6 d              | 88(3.6% from Ag-109 m), 22-26(99.4%) Ag K x-rays                                  | 50 nCi-1 mCi         | 1.85 kBq-37 MBq |
| GF-139-x       | Cerium-139    | 137.640 d            | 33.03(22.8%), 33.4(41.9%), 165.9(79.9%), 33-39(80%) La x-rays                     | 5 nCi-10 µCi         | 185 Bq-370 kBq  |
| GF-141-x       | Cerium-141    | 32.5 d               | 36.0 (9.1%), 35.6 (5%), 145.4 (48.4%), 352.42 (17%), Pr x-rays                    | On Request           | —               |
| GF-134-x       | Cesium-134    | 754.28 d             | 563(8.4%), 569(15.4%), 605(97.6%), 796(85.4%)                                     | 5 nCi-100 µCi        | 185 Bq-3.7 MBq  |
| GF-137-x       | Cesium-137    | 30.17 y              | 662(85.1% from Ba-137), 32-37(7.2%) Ba K x-rays                                   | 5 nCi-1 mCi          | 185 Bq-37 MBq   |
| GF-051-x       | Chromium-51   | 27.706 d             | 320(9.86%), 4.9-5.4(22.8%) V K x-rays   | 25 nCi-500 µCi       | 925 Bq-18.5 MBq |

### Gamma Standards—Type R

| Catalog Number | Nuclide            | Half-Life                 | Major Photon Emissions (keV)   | Available Activities |                  |
|----------------|--------------------|---------------------------|--|----------------------|------------------|
| GF-056-x       | Cobalt-56          | 77.31 d                   | 846.8(99.9%), 1238(66.1%), 1771(15.5%), 2035(7.8%), 2598(17%), 3253(7.6%), others up to 3452 | 10 nCi-100 µCi       | 370 Bq-3.7 MBq   |
| GF-057-x       | Cobalt-57          | 271.79 d                  | 14(9.2%), 122(85.6%), 136.5(10.7%), 6.4-7.1(57.9%) Fe K x-rays                               | 5 nCi-100 µCi        | 185 Bq-3.7 MBq   |
| GF-058-x       | Cobalt-58          | 70.86 d                   | 810(99.5%), 6.4-7.1(26.7%) Fe K x-rays   | 5 nCi-100 µCi        | 185 Bq-3.7 MBq   |
| GF-060-x       | Cobalt-60          | 5.272 y                   | 1173(100%), 1333(100%)   | 5 nCi-100 µCi        | 185 Bq-3.7 MBq   |
| GF-152-x       | Europium-152       | 4933 d                    | 122-1408, 40-47(74%) Sm +Gd x-rays   | 25 nCi-100 µCi       | 925 Bq-3.7 MBq   |
| GF-154-x       | Europium-154       | 3136.8 d                  | 123-1597, 42-50(25.6%) Gd x-rays   | 25 nCi-100 µCi       | 925 Bq-3.7 MBq   |
| GF-155-x       | Europium-155       | 1770 d                    | 87(34%), 105(20.6%), 42-50(24.0%), Gd K x-rays   | 10 nCi-100 µCi       | 370 Bq-3.7 MBq   |
| GF-153-x       | Gadolinium-153     | 242 d                     | 97(29.5%), 103(21.1%), 40-49(122%) Eu K x-rays   | 10 nCi-100 µCi       | 370 Bq-3.7 MBq   |
| GF-068-x       | Germanium-68       | 270.8 d                   | 511(178%), 1077(3.2%) from Ga-68, 9.2-10.4(44.1%) Ga K x-rays, 8.6-9.6(4.7%) Zn K x-rays     | 100 nCi-100 µCi      | 3.7 kBq-3.7 MBq  |
| GF-166-x       | Holmium-166m       | 1200 y                    | 81-1427, 48-58(37.6%) Er K x-rays  | 10 nCi-10 µCi        | 370 Bq-370 kBq   |
| GF-125-x       | Iodine-125         | 59.43 d                   | 35(6.58%), 27-32(139%) Te K x-rays   | 10 nCi-10 µCi        | 370 Bq-370 kBq   |
| GF-129-x       | Iodine-129         | 1.57 x 10 <sup>7</sup> y  | 40(7.5%), 29-35(70.4%) Xe K x-rays   | 50 nCi-1 µCi         | 1.85 kBq-37 kBq  |
| GF-059-x       | Iron-59            | 44.51 d                   | 1099(56.3%), 1292(43.7%)   | 5 nCi-100 µCi        | 185 Bq-3.7 MBq   |
| GF-054-x       | Manganese-54       | 312.3 d                   | 835(100%), 5.4-5.9(25.6%) Cr K x-rays  | 5 nCi-100 µCi        | 185 Bq-3.7 MBq   |
| GF-203-x       | Mercury-203        | 46.595 d                  | 279.2(81.5%)   | 10 nCi-50 µCi        | 370 Bq-1.85 MBq  |
| GF-226-x       | Radium-226         | 1600 y                    | 47-2448 (includes daughters)   | 50 nCi-10 µCi        | 1.85 kBq-370 kBq |
| GF-106-x       | Ruthenium-106      | 1.020 y                   | 512(20.7%), 622(9.8%) from Rh-106  | 25 nCi-100 µCi       | 925 Bq-3.7 MBq   |
| GF-046-x       | Scandium-46        | 83.79 d                   | 889(99.9%), 1121(99.9%)  | 5 nCi-100 µCi        | 185 Bq-3.7 MBq   |
| GF-075-x       | Selenium-75        | 119.64 d                  | 121(17.1%), 136(58.8%), 265(59%), 280(25%), 10.5-12.0(56.8%) As K x-rays                     | 5 nCi-100 µCi        | 185 Bq-3.7 MBq   |
| GF-110-x       | Silver-110m        | 249.8 d                   | 657.8(94.4%), 884.6(72.6%)   | 5 nCi-50 µCi         | 185 Bq-1.85 MBq  |
| GF-131-x       | Simulated I-131    | ~5 y                      | 356(from Ba-133), 662(from Cs-137/Ba-137)  | 50 nCi-100 µCi       | 1.85 kBq-3.7 MBq |
| GF-022-x       | Sodium-22          | 950.8 d                   | 511(178%), 1275(100%)  | 5 nCi-100 µCi        | 185 Bq-3.7 MBq   |
| GF-085-x       | Strontium-85       | 64.849 d                  | 514(98.4%), 13.3-15.3(58.7%) Rb K x-rays   | 5 nCi-100 µCi        | 185 Bq-3.7 MBq   |
| GF-228-x       | Thorium-228        | 698.2 d                   | 84-2614 (includes daughters)   | 50 nCi-10 µCi        | 1.85 kBq-370 kBq |
| GF-113-x       | Tin-113            | 115.09 d                  | 392(64% from In-113 m), 24-28(96.8%) In K x-rays   | 5 nCi-100 µCi        | 185 Bq-3.7 MBq   |
| GF-235-x       | Uranium-235        | 7.037 x 10 <sup>8</sup> y | 143(10.5%), 186(53%), 90-105(10.8%) Th K x-rays  | 10 nCi-100 nCi       | 370 Bq-3.7 kBq   |
| GF-238U-x      | Uranium (Natural)  | 4.468 x 10 <sup>9</sup> y | 26-2448 (includes daughters)   | 10 nCi-35 nCi        | 370 Bq-1.29 kBq  |
| GF-088-x       | Yttrium-88         | 106.630 d                 | 898(94%), 1836(99.4%), 14.1-16.2(61.6%) Sr K x-rays  | 5 nCi-100 µCi        | 185 Bq-3.7 MBq   |
| GF-065-x       | Zinc-65            | 244.26 d                  | 1116(50.6%), 8.0-8.9(38.7%) Cu K x-rays  | 5 nCi-100 µCi        | 185 Bq-3.7 MBq   |
| GF-095-x       | Zirconium-95/Nb-95 | 64.02 d                   | 724(44.1%), 757(54.5%)   | 10 nCi-50 µCi        | 370 Bq-1.85 MBq  |

### Gamma Sets—Type R

| Catalog Number | Available Activities | Sets Consist Of                                       |
|----------------|----------------------|---|
| GF-290-0.1x    | 0.1 µCi              | 3.7 kBq   |
| GF-290-1x      | 1 µCi                | Ba-133, Cd-109, Co-57, Co-60, Cs-137, Mn-54 and Na-22 |
| GF-290-10x     | 10 µCi               | Ba-133, Cd-109, Co-57, Co-60, Cs-137, Mn-54 and Na-22 |

# Gamma and X-Ray Standards

## Gamma Standards—Type T

The type T plastic test tube is used in clinical instrument calibrations. Each polypropylene tube contains 0.75 mL of active epoxy with the balance of the tube filled with cold epoxy. Tube sizes are 2.95" high x 0.472" diameter (75 mm x 12 mm) or 2.17" high x 0.472" diameter (55 mm x 12 mm).




| Window & Active Materials |                                     |
|---------------------------|-------------------------------------|
| Window                    | Nature of Active Materials          |
| Not Applicable            | Metallic Salts Distributed in Epoxy |

### Overall Dimensions

| X  | Height        | Diameter       | Active Volume |
|----|---------------|----------------|---------------|
| T1 | 2.17" (55 mm) | 0.472" (12 mm) | 0.75 mL       |
| T2 | 2.95" (75 mm) | 0.472" (12 mm) | 0.75 mL       |

Figure 52-A: Type T

A1901



## Gamma Standards—Type T

| Catalog Number | Nuclide       | Half-Life            | Major Photon Emissions (keV)  | Available Activities |                  |
|----------------|---------------|----------------------|---|----------------------|------------------|
| GF-241-x       | Americium-241 | 432.17 y             | 59.5 (36%), 11-20 (39.5%) Np L x-rays   | 5 nCi-10 µCi         | 185 Bq-370 kBq   |
| GF-124-x       | Antimony-124  | 60.20 d              | 602.7(97.9%), 722.8(10.9%), 1690.9(47.6%)   | 5 nCi-10 µCi         | 185 Bq-370 kBq   |
| GF-125A-x      | Antimony-125A | 1007.7 d             | 428(29.7%), 464(10.5%), 601(17.7%), 607(5%), 636(11.2%), 27-32(75.1%) Te K x-rays | 5 nCi-10 µCi         | 185 Bq-370 kBq   |
| GF-133-x       | Barium-133    | 3862 d               | 80(34.1%), 303(18.3%), 356(61.9%), 32-37(53.2%) Cs K x-rays                       | 5 nCi-10 µCi         | 185 Bq-370 kBq   |
| GF-007-x       | Beryllium-7   | On Request           |   |                      |                  |
| GF-207-x       | Bismuth-207   | $1.16 \times 10^4$ d | 570(97.7%), 1064(74.5%), 9-15(32.5%), Pb L x-rays, 72-88(77.7%) Pb K x-rays       | 5 nCi-10 µCi         | 185 Bq-370 kBq   |
| GF-109-x       | Cadmium-109   | 462.6 d              | 88(3.6% from Ag-109 m), 22-26(99.4%) Ag K x-rays                                  | 50 nCi-10 µCi        | 1.85 kBq-370 kBq |
| GF-139-x       | Cerium-139    | 137.640 d            | 33.03(22.8%), 33.4(41.9%), 165.9(79.9%), 33-39(80%) La x-rays                     | 5 nCi-10 µCi         | 185 Bq-370 kBq   |
| GF-141-x       | Cerium-141    | 32.5 d               | 36.0 (9.1%), 35.6 (5%), 145.4 (48.4%), 352.42 (17%), Pr x-rays                    | —                    | —                |
| GF-134-x       | Cesium-134    | 754.28 d             | 563(8.4%), 569(15.4%), 605(97.6%), 796(85.4%)                                     | 5 nCi-10 µCi         | 185 Bq-370 kBq   |
| GF-137-x       | Cesium-137    | 30.17 y              | 662(85.1% from Ba-137), 32-37(7.2%) Ba K x-rays                                   | 5 nCi-10 µCi         | 185 Bq-370 kBq   |
| GF-051-x       | Chromium-51   | 27.706 d             | 320(9.86%), 4.9-5.4(22.8%) V K x-rays   | 25 nCi-10 µCi        | 925 Bq-370 kBq   |

| Gamma Standards—Type T |                    |                         |  |                      |                  |  |
|------------------------|--------------------|-------------------------|--|----------------------|------------------|--|
| Catalog Number         | Nuclide            | Half-Life               | Major Photon Emissions (keV)   | Available Activities |                  |  |
| GF-056-x               | Cobalt-56          | 77.31 d                 | 846.8(99.9%), 1238(66.1%), 1771(15.5%), 2035(7.8%), 2598(17%), 3253(7.6%), others up to 3452 | 10 nCi-10 µCi        | 370 Bq-370 kBq   |  |
| GF-057-x               | Cobalt-57          | 271.79 d                | 14(9.2%), 122(85.6%), 136.5(10.7%), 6.4-7.1(57.9%) Fe K x-rays                               | 5 nCi-10 µCi         | 185 Bq-370 kBq   |  |
| GF-058-x               | Cobalt-58          | 70.86 d                 | 810(99.5%), 6.4-7.1(26.7%) Fe K x-rays   | 5 nCi-10 µCi         | 185 Bq-370 kBq   |  |
| GF-060-x               | Cobalt-60          | 5.272 y                 | 1173(100%), 1333(100%)   | 5 nCi-10 µCi         | 185B q-370 kBq   |  |
| GGF-152-x              | Europium-152       | 4933 d                  | 122-1408, 40-47(74%) Sm +Gd x-rays   | 25 nCi-10 µCi        | 925 Bq-370 kBq   |  |
| GF-154-x               | Europium-154       | 3136.8 d                | 123-1597, 42-50(25.6%) Gd x-rays   | 25 nCi-10 µCi        | 925 Bq-370 kBq   |  |
| GF-155-x               | Europium-155       | 1770 d                  | 87(34%), 105(20.6%), 42-50(24.0%), Gd K x-rays   | 10 nCi-10 µCi        | 370 Bq-370 kBq   |  |
| GF-153-x               | Gadolinium-153     | 242 d                   | 97(29.5%), 103(21.1%), 40-49(122%) Eu K x-rays   | 10 nCi-10 µCi        | 370 Bq-370 kBq   |  |
| F-068-x                | Germanium-68       | 270.8 d                 | 511(178%), 1077(3.2%) from Ga-68, 9.2-10.4(44.1%) Ga K x-rays, 8.6-9.6(4.7%) Zn K x-rays     | 100 nCi-10 µCi       | 3.7 kBq-370 kBq  |  |
| GF-166-x               | Holmium-166m       | 1200 y                  | 81-1427, 48-58(37.6%) Er K x-rays  | 10 nCi-1 µCi         | 370 Bq-37 kBq    |  |
| GF-125-x               | Iodine-125         | 59.43 d                 | 35(6.58%), 27-32(139%) Te K x-rays   | 10 nCi-10 µCi        | 370 Bq-370 kBq   |  |
| GF-129-x               | Iodine-129         | 1.5 x 10 <sup>7</sup> y | 40(7.5%), 29-35(70.4%) Xe K x-rays   | 50 nCi-1 µCi         | 1.85 kBq-37 kBq  |  |
| GF-059-x               | Iron-59            | 44.51 d                 | 1099(56.3%), 1292(43.7%)   | 5 nCi-10 µCi         | 185 Bq-370 kBq   |  |
| GF-054-x               | Manganese-54       | 312.3 d                 | 835(100%), 5.4-5.9(25.6%) Cr K x-rays  | 5 nCi-10 µCi         | 185 Bq-370 kBq   |  |
| GF-203-x               | Mercury-203        | 46.595 d                | 279.2(81.5%)   | 10 nCi-10 µCi        | 370 Bq-370 kBq   |  |
| GF-046-x               | Scandium-46        | 83.79 d                 | 889(99.9%), 1121(99.9%)  | 5 nCi-10 µCi         | 185 Bq-370 kBq   |  |
| GF-075-x               | Selenium-75        | 119.64 d                | 121(17.1%), 136(58.8%), 265(59%), 280(25%), 10.5-12.0(56.8%) As K x-rays                     | 5 nCi-10 µCi         | 185B q-370 kBq   |  |
| GF-110-x               | Silver-110m        | 249.8 d                 | 657.8(94.4%), 884.6(72.6%)   | 5 nCi-10 µCi         | 185 Bq-370 kBq   |  |
| GF-131-x               | Simulated I-131    | ~5 y                    | 356 (from Ba-133), 662 (from Cs-137/Ba-137)  | 50 nCi-10 µCi        | 1.85 kBq-370 kBq |  |
| GF-022-x               | Sodium-22          | 950.8 d                 | 511(178%), 1275(100%)  | 5 nCi-10 µCi         | 185 Bq-370 kBq   |  |
| GF-085-x               | Strontium-85       | 64.849 d                | 514(98.4%), 13.3-15.3(58.7%) Rb K x-rays   | 5 nCi-10 µCi         | 185 Bq-370 kBq   |  |
| GF-113-x               | Tin-113            | 115.09 d                | 392(64% from In-113 m), 24-28(96.8%) In K x-rays   | 5 nCi-10 µCi         | 185 Bq-370 kBq   |  |
| GF-088-x               | Yttrium-88         | 106.630 d               | 898(94%), 1836(99.4%), 14.1-16.2(61.6%) Sr K x-rays  | 5 nCi-10 µCi         | 185 Bq-370 kBq   |  |
| GF-065-x               | Zinc-65            | 244.26 d                | 1116(50.6%), 8.0-8.9(38.7%) Cu K x-rays  | 5 nCi-10 µCi         | 185 Bq-370 kBq   |  |
| GF-095-x               | Zirconium-95/Nb-95 | 64.02 d                 | 724(44.1%), 757(54.5%)   | 10 nCi-10 µCi        | 370 Bq-370 kBq   |  |

| Gamma Sets—Type T |                      |                 |   |
|-------------------|----------------------|-----------------|---|
| Catalog Number    | Available Activities | Sets Consist Of |   |
| GF-290-0.1x       | 0.1 µCi              | 3.7 kBq         | Ba-133, Cd-109, Co-57, Co-60, Cs-137, Mn-54 and Na-22 |
| GF-290-1x         | 1 µCi                | 37 kBq          | Ba-133, Cd-109, Co-57, Co-60, Cs-137, Mn-54 and Na-22 |
| GF-290-10x        | 10 µCi               | 370 kBq         | Ba-133, Cd-109, Co-57, Co-60, Cs-137, Mn-54 and Na-22 |